

MEASUREMENTS OF C_z , C_x FOR $K^+\Lambda$ AND $K^+\Sigma^0$ PHOTOPRODUCTION

ROBERT BRADFORD

*Department of Physics and Astronomy
University of Rochester
500 Wilson Boulevard
Rochester, NY 14627-0171, USA
E-mail: bradford@pas.rochester.edu*

REINHARD SCHUMACHER

*Dept of Physics
Carnegie Mellon University
5000 Forbes Avenue
Pittsburgh, PA 15213
E-mail: schumacher@cmu.edu*

FOR THE CLAS COLLABORATION.

The CLAS collaboration has recently completed first measurements of the double polarization observables C_x and C_z for the reactions $\gamma p \rightarrow K^+\Lambda$ and $\gamma p \rightarrow K^+\Sigma^0$. C_x and C_z are the beam-recoil polarization asymmetries measuring the polarization transfer from incoming circularly polarized photons to outgoing hyperons along two directions in the reaction's production plane. The Λ is found to nearly maximally polarized along the direction of incident photon's polarization for forward-going kaons. Polarization transfer to the Σ^0 is different from the Λ case.

1. Introduction

Measurement of polarization observables have long been recognized as key to unraveling baryon production mechanisms. This work represents the first measurement of C_x and C_z for $\gamma p \rightarrow K^+\Lambda$ and $\gamma p \rightarrow K^+\Sigma^0$.

2. C_x and C_z

C_x and C_z measure the polarization transfer from a circularly polarized incident photon beam to the recoiling Λ or Σ^0 baryons along two orthogonal directions in the production plane of the K^+ -hyperon system. The $\hat{x}$ and $\hat{z}$

directions are defined in the CM frame, with $\hat{z}$ lying along the directions of the incident photon beam's polarization. In this analysis, the polarization transfer was measured through the beam helicity asymmetry according to

$$A_{x/z}(\cos\theta_p) = \frac{N^+(\cos\theta_p) - N^-(\cos\theta_p)}{N^+(\cos\theta_p) + N^-(\cos\theta_p)} = \alpha_{eff}\eta C_{x/z}\cos\theta_p \quad (1)$$

where $\cos\theta_p$ is the direction of the proton from the hyperon's decay measured in the hyperon rest frame with respect to the $\hat{x}$ or $\hat{z}$ axis. η is the polarization of the incident photon beam, $N^+(\cos\theta_p)$ and $N^-(\cos\theta_p)$ are the beam helicity dependent hyperon yields in a given $\cos\theta_p$ bin. α_{eff} is the effective weak decay asymmetry parameter, and has a value of 0.642 for $K^+\Lambda$ and -0.165 for $K^+\Sigma^0$. The value of α_{eff} for the Σ^0 decay arises from our technique of measuring the proton distribution in the rest frame of the Σ^0 , not the Λ ; this dilutes its value to less than the nominal -0.642/3.


3. Experimental Setup and Analysis

The data were taken using the CLAS spectrometer in Hall B at Jefferson Lab. The experiment used a circularly polarized photon beam incident on a liquid hydrogen target. Data were taken with endpoint photon energies of 2.4 and 2.9 GeV. From this dataset, we also measured differential cross sections, which are currently available in preprint¹.


All analyzed events were required to have explicit detection of the K^+ and proton. The Λ or Σ^0 hyperons were identified in the $p(\gamma, K^+)Y$ missing mass. The data were binned in beam helicity, the cosine of the kaon angle in the CM frame ($\cos(\theta_{KCM})$), the cosine of the proton angle ($\cos(\theta_p)$) and photon energy (E_γ). Within kinematic each bin, yields were extracted by fitting a Gaussian peak to each hyperon in the missing mass spectrum. Backgrounds were modeled with a polynomial. The beam helicity asymmetry was plotted against $\cos(\theta_p)$ and the slope of this distribution was extracted with a linear fit. Complete analysis details are available elsewhere².

4. Results and Discussion

Sample results are presented in Figures 1 and 2. The data are plotted with predictions from the Kaon-MAID⁴ (solid line) and Janssen³ (dashed line) isobar models. The data plotted are for only a few representative bins in $\cos(\theta_{KCM})$. The full results include nine bins in $\cos(\theta_{KCM})$ for $K^+\Lambda$ and six bins for $K^+\Sigma^0$.



(a)



(b)

Figure 1. C_x and C_z for $K^+\Lambda$ in two different kaon-angle bins. Top: $\cos(\theta_K) = -0.75$, bottom: $\cos(\theta_K) = 0.25$. The data are a subset of the 2005 CLAS results. The curves are predictions from the Kaon-MAID (solid line,⁴) and Janssen (dashed line,³) isobar models.




Figure 2. C_x and C_z for $K^+\Sigma^0$ for $\cos(\theta_K) = 0.5$. The data are a subset the 2005 CLAS results. The curves are predictions from the Kaon-MAID (solid line, ⁴) and Janssen (dashed line, ³) isobar models.

The Λ results show some W -dependent structure at backward kaon angles and then stabilize at more forward-going kaon angles. For this hyperon, C_z is near one over most of the forward hemisphere of the kaon angle while C_x is near zero for the same range.

The $K^+\Sigma^0$ results show no preferred direction for the polarization over the kaon angle range. The precision of these results here appears worse due to the small value of α_{eff} .

Of the models shown, the Janssen does a good job of following the data. The MAID curve does not fair as well. This model has the oddity of predicting that C_z saturates at -1 in $K^+\Lambda$ for forward-going kaons.

References

1. R. Bradford, *et al.* Preprint available at: nucl-ex/0509033. Submitted to *Phys. Rev. C*.
2. R. Bradford, *Measurement of differential cross sections and C_x and C_z for $\gamma p \rightarrow K^+\Lambda$ and $\gamma p \rightarrow K^+\Sigma^0$ using CLAS at Jefferson Lab*. Ph.D. thesis, Carnegie Mellon University, 2005, to be published. Available on-line at: <http://www.jlab.org/Hall-B/general/thesis/bradford/index.html>
3. S. Janssen, *Strangeness production on the nucleon*. Ph.D. thesis, University of Gent, 2002.
4. F.X. Lee, *et al.*, *Nucl. Phys. A* **695**, 237 (2001).